From Cough to Cure: A Path of Ideal Behaviors in Tuberculosis Control

B A R R I E R S

LEVELS	* Poor knowledge of TB symptoms	* Misperception of costs	* Poor knowledge of diagnostic steps	* Poor knowledge of length of treatment	* Poor knowledge of length of treatment	* Poor knowledge of length of treatment
ndividual & G roup	 * Poor knowledge of TB care and cure * Stigma related to TB diagnosis * Low risk preception 	 * Preference for non- DOTS health services * Attitude about health services * Stigma * Social Norm 	* Expectations about medical services (get meds not tests)	* Stigma	* Stigma	* Stigma
IDEAL BEHAVIOR	Seek Care	Go to DOTS	Complete Diagnosis	Begin Treatment	Continue Treatment & follow-up smears	Complete Treatment & final smear
S ystem		 * Time, cost, distance to DOTS facility * Lack of linkages between DOTS and other providers (non- DOTs & HIV care) * Missed diagnosis and / or lack of referral by non-DOTS providers 	 * Providers' poor knowledge of correct procedures * Providers' poor inter-personal communication * Lack of resources, including human resources * Poor quality of services (hours, wait- time) 	 * Time, cost, distance to DOTS facility * Poor quality of services * Health providers fail to give adequate information * Lack of medications 	 * Time, cost, distance to DOTS facility * Poor quality of services * Health providers fail to give adequate information * Lack of medications 	 * Time, cost, distance to DOTS facility * Poor quality of services * Health providers fail to give adequate information * Lack of medications
			time) * Time, cost and distance			